arXiv:0808.1829v1 [math.NT] 13 Aug 2008

NOTE ON THE EULER NUMBERS AND POLYNOMIALS

Taekyun Kim

Abstract. In this paper we investigate the properties of the Euler functions. By using the Fourier transform for the Euler function, we derive the interesting formula related to the infinite series. Finally we give some interesting identities between the Euler numbers and the second kind stirling numbers.

§1. Introduction/Definition

The constants E_k in the Taylor series expansion

$$F(t)=\frac{2}{e^t+1}=\sum_{n=0}^{\infty}E_n\frac{t^n}{n!},\ \text{for } |t|\leq\pi,\ \text{(see [1-31])}, \tag{1}$$

are known as the Euler numbers. The first few are 1, $-\frac{1}{2}$, 0, $\frac{1}{4}, \cdots$, and $E_{2k}=0$ for $k=1,2,3,\cdots$. These numbers arise in the series expansions of trigonometric functions, and are extremely important in number theory and analysis. The Euler polynomials, $E_n(x)$, are defined as

$$F(x,t)=F(t)e^{xt}=\frac{2}{e^t+1}e^{xt}=\sum_{n=0}^{\infty}E_n(x)\frac{t^n}{n!},\ \text{for } x\in\mathbb{R}. \tag{2}$$

For $x\in\mathbb{R}$ with $0\leq x<1$, the Euler polynomials are called the Euler functions. From (1) and (2) we can derive

$$E_n(x)=\sum_{l=0}^{n}\binom{n}{l}E_lx^{n-l},\ \text{where } \binom{n}{l}=\frac{n(n-1)\cdots(n-l+1)}{l!}. \tag{3}$$

Key words and phrases. Euler numbers,Fourier transform, infinite series, Euler function.
2000 AMS Subject Classification: 11B68, 11S80

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

Thus, we obtain the distribution relation for the Euler polynomials as follows. For $d \in \mathbb{N}$ with $d \equiv 1 (\mod 2)$, we have

$$\sum_{k=0}^{d-1} (-1)^k E_n(\frac{x+k}{d}) = d^{-n} E_n(x).$$

By (1), it is easy to see that the recurrence relation for the Euler numbers is given by

$$E_0 = 1, \text{ and, } \sum_{l=0}^{n} \binom{n}{l} E_l + E_n = 2\delta_{0,n} \text{ where } \delta_{0,n} \text{ is Kronecker symbol.} \tag{4}$$

From (3) and (4), we note that

$$E_n(1) = \sum_{l=0}^{n} \binom{n}{l} E_l = -E_n, \text{ for } n \geq 1. \tag{5}$$

Thus, we obtain the following lemma.

Lemma 1. *For $n \in \mathbb{N}$, we have $E_n(1) = -E_n$.*

From (3) we can easily derive

$$\begin{aligned} \frac{dE_n(x)}{dx} &= \frac{d}{dx} \sum_{k=0}^{n} \binom{n}{k} E_k x^{n-k} = \sum_{k=0}^{n} \binom{n}{k} (n-k) E_k x^{n-k-1} \\ &= n \sum_{k=0}^{n} \frac{(n-1)!}{(n-k-1)!k!} E_k x^{n-1-k} = n \sum_{k=0}^{n-1} \binom{n-1}{k} E_k x^{n-1-k} = n E_{n-1}(x). \end{aligned} \tag{6}$$

By (6), we obtain the following proposition.

Proposition 2. *For $n \geq 0$, we have*

$$\int_0^x E_n(t)dt = \frac{1}{n+1} E_{n+1}(x). \tag{7}$$

In this paper we investigate the properties of the Euler functions. By using the Fourier transform for the Euler function, we derive the interesting formula related to the infinite series. Finally we give some interesting identities between the Euler numbers and the second kind stirling numbers.

§2. Euler Functions

In this section, we assume that $E_n(x)$ is the Euler function. Let us consider the Fourier transform for the Euler function, $E_n(x)$, as follow. For $m \in \mathbb{N}$, the Fourier transform on the Euler function is given by

$$E_m(x) = \sum_{n=-\infty}^{\infty} a_n^{(m)} e^{(2n+1)\pi ix}, \ (a_n^{(m)} \in \mathbb{C}), \tag{8}$$

where

$$a_n^{(m)} = \int_0^1 E_m(x) e^{-(2n+1)\pi ix} dx. \tag{9}$$

From (9), we note that

$$\begin{aligned} a_n^{(m)} &= \int_0^1 E_m(x) e^{-\pi i(2n+1)x} dx \\ &= \left[\frac{E_{m+1}(x)}{m+1} e^{-\pi i(2n+1)x} \right]_0^1 + \frac{(2n+1)\pi i}{m+1} \int_0^1 E_{m+1}(x) e^{-(2n+1)\pi ix} dx \\ &= \frac{(2n+1)\pi i}{m+1} \int_0^1 E_{m+1}(x) e^{-(2n+1)\pi ix} dx = \frac{(2n+1)\pi i}{m+1} a_n^{(m+1)}. \end{aligned} \tag{10}$$

Thus, we have

$$a_n^{(m)} = \frac{m}{(2n+1)\pi i} a_n^{(m-1)} = \frac{m(m-1)}{((2n+1)\pi i)^2} a_n^{(m-2)} = \cdots = \frac{m!}{((2n+1)\pi i)^{m-1}} a_n^{(1)}. \tag{11}$$

It is easy to see that

$$\begin{aligned} a_n^{(1)} &= \int_0^1 E_1(x) e^{-(2n+1)\pi ix} dx = \int_0^1 (x - \frac{1}{2}) e^{-(2n+1)\pi ix} dx \\ &= -\frac{1}{(2n+1)\pi i} \left[(x - \frac{1}{2}) e^{-(2n+1)\pi ix} \right]_0^1 + \frac{1}{(2n+1)\pi i} \int_0^1 e^{-(2n+1)\pi ix} dx \\ &= \frac{2}{((2n+1)\pi i)^2}. \end{aligned} \tag{12}$$

From (11) and (13), we can derive

$$a_n^{(m)} = 2 \frac{m!}{((2n+1)\pi i)^{m+1}}, \ m \in \mathbb{N}, \text{ and } a_n^{(0)} = \frac{2}{(2n+1)\pi i}. \tag{13}$$

By (8) and (13), we see that

$$E_m(x) = m! 2 \sum_{n=-\infty}^{\infty} \frac{e^{(2n+1)\pi ix}}{((2n+1)\pi i)^{m+1}}, \text{ for } 0 \le x < 1.$$

Therefore, we obtain the following theorem.

Theorem 3. *For $m \in \mathbb{Z}_+ = \mathbb{N} \bigcup \{0\}$, $x \in \mathbb{R}$ with $0 \leq x < 1$, we have*

$$E_m(x) = m!2 \sum_{n=-\infty}^{\infty} \frac{e^{(2n+1)\pi i x}}{((2n+1)\pi i)^{m+1}}.$$

If we take $x = 1$, then we have

$$E_m(1) = -m!2 \sum_{n=-\infty}^{\infty} \frac{1}{((2n+1)\pi i)^{m+1}}. \tag{14}$$

By(14) and Lemma 1, we obtain the following corollary.

Corollary 4. *For $m \in \mathbb{Z}_+ = \mathbb{N} \bigcup \{0\}$, we have*

$$E_m = m!2 \sum_{n=-\infty}^{\infty} \frac{1}{((2n+1)\pi i)^{m+1}}.$$

From Corollary 4, we note that

$$E_{2m+1} = (-1)^{m+1} 2 \frac{(2m+1)!}{\pi^{2m+2}} \sum_{n=-\infty}^{\infty} \frac{1}{(2n+1)^{2m+2}}.$$

Thus, we have

$$\sum_{n=-\infty}^{\infty} \frac{1}{(2n+1)^{2m+2}} = (-1)^{m+1} \frac{E_{2m+1}}{2(2m+1)!} \pi^{2m+2}. \tag{15}$$

By (15), we obtain the following corollary.

Corollary 5. *For $m \in \mathbb{Z}_+$, we have*

$$\sum_{n=1}^{\infty} \frac{1}{(2n+1)^{2m+2}} = (-1)^{m+1} \frac{E_{2m+1}}{4(2m+1)!} \pi^{2m+2}.$$

Note that

$$\begin{aligned} \frac{1}{1+e^{-x}} &= \sum_{n=0}^{\infty} e^{-nx}(-1)^n = \sum_{n=0}^{\infty} (e^{-x})^n (-1)^n = \sum_{n=0}^{\infty} \left(\sum_{k=0}^{\infty} \frac{(-1)^k}{k!} x^k \right)^n (-1)^n \\ &= \sum_{n=0}^{\infty} (-1)^n \left(\sum_{a_1+a_2+\cdots=n} \frac{n!}{a_1!a_2!\cdots} \frac{(-1)^{a_1+2a_2+\cdots}}{(1!)^{a_1}(2!)^{a_2}\cdots} \right) x^{a_1+2a_2+\cdots}. \end{aligned} \tag{16}$$

Let $p(i,j) : a_1 + 2a_2 + \cdots = i, a_1 + a_2 + \cdots = j$. From (16), we note that

$$\frac{1}{1+e^{-x}} = \sum_{m=0}^{\infty} \left(\sum_{n=0}^{m} (-1)^n \sum_{p(m,n)} \frac{n!}{a_1!a_2!\cdots a_m!} \frac{(-1)^m x^m}{(1!)^{a_1}\cdots(m!)^{a_m}} \right)$$

$$(17) \qquad = \sum_{m=0}^{\infty} (-1)^m \left(\sum_{n=0}^{m} n!(-1)^n \sum_{p(m,n)} \frac{m!}{a_1!a_2!\cdots a_m!} \frac{(-1)^m}{(1!)^{a_1}(2!)^{a_2}\cdots(m!)^{a_m}} \right) \frac{x^m}{m!}$$

$$= \sum_{m=0}^{\infty} (-1)^m \sum_{n=0}^{m} n!(-1)^n s_2(m,n) \frac{x^m}{m!},$$

where $s_2(m,n)$ is the second kind stirling number.

By the definition of Euler number, we easily see that

$$(18) \qquad \frac{1}{1+e^{-x}} = \frac{1}{2}\left(\frac{2}{1+e^{-x}}\right) = \frac{1}{2}\sum_{m=0}^{\infty}(-1)^m E_m \frac{x^m}{m!}, \text{ (see [5]).}$$

By comparing the coefficients of $\frac{t^n}{n!}$ on the both sides of (17) and (18), we obtain

$$E_m = 2\sum_{n=0}^{\infty}(-1)^n n! s_2(m,n),$$

where $s_2(m,n)$ is the second kind stirling number.

References

1. M. Cenkci, Y. Simsek, V. Kurt, *Further remarks on multiple p-adic q-L-function of two variables*, Adv. Stud. Contemp. Math. **14** (2007), 49-68.
2. M. Cenkci, *The p-adic generalized twisted (h,q)-Euler-l-function and its applications*, Adv. Stud. Contemp. Math. **15** (2007), 37-47.
3. M. Cenkci, M. Can, *Some results on q-analogue of the Lerch zeta function*, Adv. Stud. Contemp. Math. **12** (2006), 213-223.
4. M. Cenkci, M. Can, V. Kurt, *p-adic interpolation functions and Kummer-type congruences for q-twisted and q-generalized twisted Euler numbers*, Adv. Stud. Contemp. Math. **9** (2004), 203-216.
5. T. Kim, *Euler Numbers and Polynomials Associated with Zeta Functions*, Abstract and Applied Analysis **2008** (2008), Article ID 581582, 11 pages.
6. T. Kim, *q-Euler numbers and polynomials associated with p-adic q-integrals*, Journal of Nonlinear Mathematical Physics **14** (2007), 15-27.
7. T. Kim, *q-Volkenborn Integration*, Russian J. Math. Phys. **9** (2002), 288-299.
8. T. Kim, *q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russian J. Math. Phys. **15** (2008), 51-57.

9. T. Kim, *q-extension of the Euler formula and trigonometric functions*, Russian J. Math. Phys. **14** (2007), 275-278.

10. T. Kim, J. Y. Choi, J. Y. Sug, *Extended q-Euler numbers and polynomials associated with fermionic p-adic q-integral on* $\mathbb{Z}_p$, Russian J. Math. Phys. **14** (2007), 160-163.

11. T. Kim, *q-generalized Euler numbers and polynomials*, Russian J. Math. Phys. **13** (2006), 293-298.

12. T. Kim, *Multiple p-adic L-function*, Russian J. Math. Phys. **13** (2006), 151-157.

13. T. Kim, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russian J. Math. Phys. **12** (2005), 186-196.

14. T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russian J. Math. Phys. **11** (2004), 71-76.

15. T. Kim, *On Euler-Barnes multiple zeta functions*, Russian J. Math. Phys. **10** (2003), 261-267.

16. T. Kim, *The modified q-Euler numbers and polynomials*, Adv. Stud. Contemp. Math. **16** (2008), 161-170.

17. T. Kim, *A note on p-adic q-integral on* $\mathbb{Z}_p$ *associated with q-Euler numbers*, Adv. Stud. Contemp. Math. **15** (2007), 133-137.

18. T. Kim, *A note on p-adic invariant integral in the rings of p-adic integers*, Adv. Stud. Contemp. Math. **13** (2006), 95-99.

19. H. Ozden, Y. Simsek, S.-H. Rim, I. N. Cangul, *On interpolation functions of the twisted generalized Frobenius-Euler numbers*, Adv. Stud. Contemp. Math. **15** (2007), 187-194.

20. H. Ozden, Y. Simsek, I. N. Cangul, *Multivariate interpolation functions of higher-order q-Euler numbers and their applications*, Abstract and Applied Analysis **2008** (2008), Art. ID 390857, 16 pages.

21. H. Ozden, Y. Simsek, I. N. Cangul, *Euler polynomials associated with p-adic q-Euler measure*, General Mathematics **15** (2007), 24-37.

22. Y. Simsek, *Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions*, Adv. Stud. Contemp. Math. **16** (2008), 251-278.

23. Y. Simsek, Y. Osman, V. Kurt, *On interpolation functions of the twisted generalized Frobenius-Euler numbers*, Adv. Stud. Contemp. Math. **15** (2007), 187-194.

24. Y. Simsek, *Hardy character sums related to Eisenstein series and theta functions*, Adv. Stud. Contemp. Math. **12** (2006), 39-53.

25. Y. Simsek, *Remarks on reciprocity laws of the Dedekind and Hardy sums*, Adv. Stud. Contemp. Math. **12** (2006), 237-246.

26. Y. Simsek, *Theorems on twisted L-function and twisted Bernoulli numbers*, Adv. Stud. Contemp. Math. **11** (2005), 205-218.

27. Y. Simsek, D. Kim, S.-H. Rim, *On the two-variable Dirichlet q-L-series*, Adv. Stud. Contemp.

Math. **10** (2005), 131-142.

28. Y. Simsek, A. Mehmet, *Remarks on Dedekind eta function, theta functions and Eisenstein series under the Hecke operators*, Adv. Stud. Contemp. Math. **10** (2005), 15-24.
29. Y. Simsek, Y. Sheldon, *Transformation of four Titchmarsh-type infinite integrals and generalized Dedekind sums associated with Lambert series*, Adv. Stud. Contemp. Math. **9** (2004), 195-202.
30. Y. Simsek, *On p-adic twisted q-L-functions related to generalized twisted Bernoulli numbers*, Russian J. Math. Phys. **13** (2006), 340-348.
31. T. Kim, *q-Riemann zeta function*, International Journal of Mathematics and Mathematical Sciences **2004, no.12** (2004), 509-605.

Taekyun Kim
Division of General Education-Mathematics,
Kwangwoon University, Seoul, S. Korea
e-mail: tkkim@kw.ac.kr; tkim64@hanmail.net